

## ORIGINAL ARTICLES

# Correlation of clinical score to pulmonary function and oxygen saturation in children with asthma attack

Z. Kaya and I. Turktaş

<sup>a</sup>Division of Allergy and Asthma. Departments of Pediatrics. Medical Faculty of Gazi University. Beşevler. Ankara. Turkey.

---

### ABSTRACT

*Background:* The aim of this study is to demonstrate the importance of the relation between clinical score, pulse oximetry and spirometric tests in an asthma attack.

*Methods:* In this randomized, double blind, observational study, 110 children (age 2-15 years) with an asthma attack who were admitted to emergency room were evaluated. Patient history, physical examination, clinical score and oxygen saturation were recorded in all patients; however pulmonary function tests were obtained only in 54 children who were over 5 years of age. The clinical score was derived from respiratory rate, wheezing, dyspnea and retractions.

*Results:* Both oxygen saturation and spirometric tests were found to be significantly correlated with the clinical score in children.

*Conclusion:* The clinical score could be used for assessing the severity of the asthma attack particularly in developing countries where laboratory facilities are not available or pulmonary function tests are not feasible.

**Key words:** Clinical score. Pulse oximetry. Spirometric tests. Asthma attack.

---

### INTRODUCTION

Acute asthma is the most common medical emergency in children and is responsible for increased hospitalization and death rates in several developing countries. In the emergency room setting, the most critical issue facing the attending physician is, deciding the adequacy of outpatient therapy of acute asthma, or when hospitalization is indicated. Traditionally, this decision is used to be made according to the clinical history, physical examination, laboratory results and response to therapy of the patient<sup>1-6</sup>. Several clinical scoring systems are presented for simplification of this. Among these some are not feasible as they contain parameters (pulsus paradoxus or inspiratory:expiratory ratio) which are difficult to be assessed by other personnel rather than the physician particularly in young children or those require active use of a monitoring device such as peak flow meter or oxygen saturation<sup>7,8</sup>. However, the clinical score we used is derived from the observation of physical signs by different care givers (physicians, nurses, respiratory therapists) in asthmatic children (table I)<sup>9</sup>.

Pulse oximetry on the other hand, is a widely used, noninvasive instrument for monitoring oxygen saturation. Its ease of performance has led to its widespread use in intensive care units and in the management of a variety of respiratory disorders<sup>7</sup>. Although obtaining spirometric test might be difficult in infants and young children who are excessively active, irritable, or uncooperative the forced expiratory volume in

Correspondence:

Zühre Kaya  
Birlik mahallesi 68. Sokak No: 18/4  
06610 Çankaya. Ankara. Turkey.  
Phone number: 90 0312 4954272  
E-mail 1: kayaz74@mynet.com  
E-mail 2: zuhrekaya@gazi.edu.tr

**Table I**  
**Clinical score<sup>9</sup>**

	0 Point	1 Point	2 Point	3 Point
<i>Respiratory rate</i> (Breath/min)				
2-3 yrs	—	≤ 34	35-39	≥ 40
4-5 yrs		≤ 30	31-35	≥ 36
6-12 yrs		≤ 26	27-30	≥ 31
> 12 yrs		≤ 23	24-27	≥ 28
<i>Wheezing</i>	None	End-expiratory wheezing	Expiratory Wheezing	Insp + Exp wheezing, or diminished breath sound or both
<i>Retractions</i>	None	Intercostal	Intercostal + Substernal	Intercostal, Substernal, + Supraclavicular
<i>Dyspnea</i>				
2-4 yrs	Normal feeding, vocalizations, and activity	1 of the following: Decreased appetite, increased coughing after play	2 of the following: Decreased appetite, increased coughing after play	Stop eating or drinking, stops playing, or drowsy or confused
≥ 5 yrs	Counts to ≥ 10 in one breath	Counts to 7-9 in one breath	Counts to 4-6 in one breath	Counts to ≤ 3 in one breath

1 sec (FEV<sub>1</sub>) correlates well with the severity of obstructive diseases in children over 5 years of age<sup>8</sup>.

The purpose of this study was to investigate the value of clinical signs in children with acute asthma and whether the information obtained would be of particular help to physicians who do not have ease of access to pulmonary function and oxygen saturation (SpO<sub>2</sub>) testing in the evaluation of such patients.

**MATERIALS AND METHODS**

We conducted a randomized double blind, observational study in 110 children (2-15 years of age) who were admitted to the Pediatric Emergency Room in more than one attack of acute bronchial asthma requiring bronchodilator therapy within 6 months of entry in the study. Children with a significant acute or chronic disease such as pneumonia, croup, pertussis, foreign body aspiration and cardiac congenital malformations were excluded. The study was approved by the Ethics Committee, and informed written consent from the parents was obtained.

The severity of acute asthma attack was evaluated by a physician using the clinical score which included respiratory rate, wheezing, dyspnea and retractions with the child standing, not crying, without fever and breathing room air only (table I)<sup>9</sup>. Simulta-

neously, a nurse, who was unaware of the patient's clinical score, measured SpO<sub>2</sub> with a pulse oximeter (Kontron Minimon 7138 Plus) using a pediatric probe placed on the finger. The mean SpO<sub>2</sub> was recorded after three measurements. Pulmonary function test (especially FEV<sub>1</sub>) with a spirometer (MIR, Sensor Medics 2130, USA) was only performed in children over 5 years of age. We recorded gender and age and measured height and weight of each children before spirometry. We have used Morris Polgar reference oxymetry values for height and weight.

Treatment was planned according to the emergency room asthma guidelines (GINA) and simply consisted of oxygen and steroid besides nebulized salbutamol<sup>10</sup>. We have also treated moderate and severe attack of asthma patients with oxygen, bronchodilator and steroid therapy. Patients were reevaluated at hourly intervals by the same clinical score, SpO<sub>2</sub> and FEV<sub>1</sub>. The maximal duration of patients in the emergency room was 6 hours and all patients with a clinical score over 7 (n: 44) on admission needed hospitalization.

Data were analyzed by SPSS-10 program and the results were shown as mean and standard deviation (mean ± SD). The relation between the clinical score, FEV<sub>1</sub> and SpO<sub>2</sub> were assessed by Pearson correlation analysis and p < 0.05 was accepted as the significance level.

**Table II**  
**Mean SpO<sub>2</sub> and FEV<sub>1</sub> values by clinical score and age group**

Clinical score	SpO <sub>2</sub>						FEV <sub>1</sub>	
	Total		2-5 year		6-15 year		6-15 year	
	X ± SD	n	X ± SD	n	X ± SD	n	X ± SD	n
0-3	92 ± 1.5	18	92 ± 1.3	7	92 ± 1.5	11	70 ± 1.9	11
4-7	90 ± 1.3	74	90 ± 1.2	41	89 ± 1.3	33	59 ± 1.8	33
8-12	88 ± 1.2	18	88 ± 1.3	8	88 ± 1.2	10	46 ± 3.5	10

X ± SD\*: mean ± standard deviation.  
n\*: patient number.

## RESULTS

110 patients with acute asthma attack (66 male, 44 female) with a median age of 8 years (range 2-15 years) were included in the study. Mean clinical score was 6 ± 2 points, mean SpO<sub>2</sub> value was 90 ± 2 %. FEV<sub>1</sub> measurement was found to be 60 ± 12.3 % in patients 5 years and older (n: 54) (table II).

FEV<sub>1</sub> measurement at presentation significantly correlated to the clinical score. Patients with clinical scores over 7 (severe clinical disease) had significantly lower FEV<sub>1</sub> values. Disruption of pulmonary function most significantly correlated to the severity of retractions followed by the severity of dyspnea and wheezing whereas SpO<sub>2</sub> value on presentation correlated most closely with the overall clinical score (table III, IV). All patients with a clinical score over 7 had a SpO<sub>2</sub> value below 91 % however of the individual clinical variables, the most significant correlation was also between the retraction severities to SpO<sub>2</sub>.

Finally, a positive correlation was also found between SpO<sub>2</sub> and FEV<sub>1</sub> (r = 0.49, p < 0.001) most significantly in patients with SpO<sub>2</sub> values below 91 %, and FEV<sub>1</sub> value below 70 % that might indicate hypoxemia.

## DISCUSSION

This study aimed to investigate the contribution of clinical evaluation in deciding hospitalization of children with acute asthma.

Previous studies demonstrated that there was no correlation between the clinical score of the children on presentation and the severity of hypoxia (r = -0.14)<sup>11,12</sup>. Meanwhile, many clinical scores have

**Table III**

**Correlation (r) between clinical score and SpO<sub>2</sub> and FEV<sub>1</sub> values in all patients**

Variable	FEV <sub>1</sub> (n: 54)	SpO <sub>2</sub> (n: 110)
Clinical score	-0.64	-0.69
Retractions	-0.48	-0.65
Wheezing	-0.12	-0.47
Respiratory rate	-0.27	-0.58
Dyspnea	-0.46	-0.46

**Table IV**

**Correlation between clinical score and SpO<sub>2</sub> values for ages**

Variable	Under 5 years of age (SpO <sub>2</sub> ) (r) (n:56)	Over 5 years of age (SpO <sub>2</sub> ) (r) (n:54)
Clinical score	-0.77	-0.69
Retractions	-0.91	-0.90
Wheezing	-0.48	-0.40
Respiratory rate	-0.70	-0.61
Dyspnea	-0.80	-0.76

been developed to evaluate hospitalized children with asthma or bronchiolitis. The one that we have chosen is easy as it could be used either by a physician or a nonphysician provider<sup>9</sup>. Using this scoring system we found a significant correlation between the clinical score and FEV<sub>1</sub> in this study. In adults and children, use of accessory respiratory muscles is reported to be the most important clinical sign and the best to correlate with the disease severity<sup>13,14</sup>. Our results also reveal that pulmonary function deteriora-

tion is found to be significantly correlated to the degree of accessory muscle use followed by the degree of dyspnea and wheezing. However, Becker et al<sup>7</sup> showed a good correlation between pulmonary function testing and pulmonary index score which was 7.3 in patients requiring admission to the hospital. Kerem et al<sup>8</sup> reported that patients who have accessory muscle use, wheezing and dyspnea had a negative correlation to FEV<sub>1</sub> values that are found to be  $\leq 60\%$  of predicted. In spite of this strong correlation, measurement of FEV<sub>1</sub> could not be helpful enough in very young, very ill or uncooperative patients besides in asthmatic children who had not previously had this test performed. On the other hand, as airway obstruction and hypoxemia could exist in the absence of clinical signs; pulmonary function tests must be used in conjunction with careful clinical evaluation.

This study depicted that a clinical score over 7 points had a sensitivity of 81 %, specificity of 100 % and positive predictive value of 100 % in diagnosing hypoxemia (SpO<sub>2</sub> < 91 %). Although there is no definite value for defining hypoxemia according to SpO<sub>2</sub> it is considered to be below 90-92 % in recent reports<sup>12,15,16</sup>.

There was also a significant correlation ( $r = -0.72$ ;  $p < 0.0001$ ) between the clinical score and SpO<sub>2</sub> in our patients. Significant correlations between clinical score and SpO<sub>2</sub> in patients under 5 years of age were similar to the patients over 5 years of age of whom spirometric tests were also performed (under 5 years  $r = -0.77$ , up to 5 years  $r = -0.69$ ). At the end of two hours period in emergency room, 44 children required hospitalization and of these 36 were under 5 years old. Patients under 5 years of age often require hospitalization as a number of anatomic and physiologic variations of early life predispose to obstructive airway disease, the symptoms progress rapidly and noticed easily<sup>17</sup>. Patients who subsequently required admission to the hospital was found to have a mean clinical score greater than 7 and SpO<sub>2</sub> lower than 91 % before the initial treatment in the emergency room in our study. Sole et al<sup>16</sup> found a significant negative correlation between the clinical scores and SpO<sub>2</sub> following emergency treatment. In addition, they showed that SpO<sub>2</sub> levels  $\leq 92\%$  were associated with a 6.3-fold greater risk for requiring additional treatment. A single clinical finding will not be very indicative of severity of airway obstruction, but a combination of findings may provide more information to the first evaluation of acute asthma. Many studies showed that the clinical score could be more beneficial when it is taken as a whole, instead of evaluating each finding alone. Kerem et al<sup>8</sup> also reported that all patients found to

have a negative correlation between accessory muscle use, respiratory rate, dyspnea and SpO<sub>2</sub> on presentation. Connett et al<sup>12</sup> showed a good correlation ( $r = -0.37$ ,  $p < 0.01$ ) in 75 hospitalized asthmatic children (mean age 3.3 years; range 1.5-14.5 years) between the asthma severity score and SpO<sub>2</sub> at the beginning of the treatment. They also demonstrated that a cut off value of SpO<sub>2</sub> below 91 % in room air was a good predictor for hospitalization in children after pre and postnebulizer treatment of acute asthma.

Many developing countries do not have pulse oximetry and spirometry to determine the need to additional intensive unit therapy. We believe that clinical score is a valuable clinical tool that could easily be used by nursing and medical staff when these are not available.

In conclusion, although the use of clinical score should not replace laboratory evaluation such as determination of oxygen saturation and pulmonary function tests in an asthma attack, we recommend starting more intensive therapy and closer supervision after hospital admission in children with asthma attack who have a clinical score over 7 especially in developing countries with inadequate sources.

## REFERENCES

1. Murray CJ, Lopez AD. Mortality by cause for eight regions of the world: Global Burden of Disease Study. *Lancet*. 1997;349:1269-76.
2. Weiss KB, Wagener DK. Changing patterns of asthma mortality: Identifying target populations at high risk. *JAMA*. 1990;264:1683-7.
3. Buist AS, Vollmer WM. Reflections on the rise in asthma morbidity and mortality. *JAMA*. 1990;264:1719-20.
4. Storr J, Barrell E, Lenney W. Rising asthma admissions and self referral. *Arch Dis Child*. 1988;63:774-9.
5. Anderson HR. Increase in hospital admissions for childhood asthma: trends in referral severity, and readmissions from 1970 to 1985 in a health region of the United Kingdom. *Thorax*. 1989;44:614-9.
6. Carman PG, Landau LI. Increased pediatric admissions with asthma in Western Australia – a problem of diagnosis? *Med J Aust*. 1990;152:23-6.
7. Becker AB, Nelson NA, Simons FE. The Pulmonary index. *AJDC*. 1984;138:574-6.
8. Kerem E, Canny G, Tibshirani R, Reisman J, Bentur L, Schuh S, Levison H. Clinical-physiologic correlations in acute asthma of childhood. *Pediatrics*. 1991;87:481-6.
9. Liu LL, Gallaher MM, Davis RL, Rutter CM, Lewis TC, Marcuse EK. Use of a respiratory clinical score among different providers. *Pediatr Pulmonol*. 2004;37:243-8.
10. National Heart, Lung and Blood Institute (1998). Pocket Guide for Asthma Management and Prevention. NIH Publication No:96-3659B, Bethesda, MD, USA.
11. Hurwitz ME, Burney RE, Howatt WF, Crowley D, Mackenzie JR. Clinical scoring does not accurately assess hypoxemia in pediatric asthma patients. *Ann Emerg Med*. 1984;13:1040-3.

12. Alario A, Lewander W, Dennehehy P, Seifer R, Mansell A. The relation between oxygen saturation and the clinical assessment of acutely wheezing infants and children. *Pediatr Emerg Care*. 1995;11:331-9.
13. Connett GJ, Lenney W. Use of pulse oximetry in the hospital management of acute asthma in childhood. *Pediatr Pulmonol*. 1993;15:345-9.
14. Margolis PA, Ferkol TW, Marsocci S, Super DM, Keyes LL, Mcnutt R, Harrell FE. Accuracy of the clinical examination in detecting hypoxemia in infants with respiratory illness. *J Pediatr*. 1994;124:552-60.
15. Geelhoed G, Landau L, Le Souef P. Evaluation of SpO<sub>2</sub> as a predictor of outcome in 280 children presenting with acute asthma. *Ann Emerg Med*. 1994;23:1236-41.
16. Sole D, Komatsu MK, Carvalho KV, Naspitz CK. Pulse oximetry in the evaluation of the severity of acute asthma and/or wheezing in children. *J Asthma*. 1999;36:327-33.
17. Martinez FD, Wright AL, Taussig LM, Holberg CJ, Halonen M, Morgan WJ. Asthma and wheezing in the first six years of life. The Group Health Medical Associates. *N Engl J Med*. 1995;332:133-8.