



ORIGINAL ARTICLE

Subjective method of refractometry and depth of focus

Nikolai M. Sergienko, Anastasia Gromova, Nikolai Sergienko*

Department of Ophthalmology, National Medical Academy of Postgraduate Education, Kiev, Ukraine

Received 1 September 2011; accepted 12 February 2012

Available online 5 April 2012

KEYWORDS

Refractometry;
Depth of field;
Depth of focus;
Pseudoaccommodation

Abstract

Purpose: To study the impact of the depth of focus on subjective refraction and distribution of myopic and hyperopic refractions.

Methods: A total of 450 eyes of 305 subjects in the age range of 23–34 years were recruited for the study. A distribution of refractions was examined using a traditional method of the results was made on the assumption that the emmetropic eye retains high visual acuity when applying convex lenses with values which are fewer or equal to the depth of focus values. The following values of the depth of focus were used: ± 0.55 D, ± 0.35 D and ± 0.2 D for visual acuity 1.0, 1.5 and 2.0, respectively.

Results: Application of the traditional method of refractometry produced the following occurrence of refractions: hypermetropia 59.3%, myopia 22% and emmetropia 18.7%. After correction of the initial results of values of the depth of focus the distribution of refractions was as follows: hypermetropia 12.7%, myopia 22% and emmetropia 65.3%.

Conclusion: The traditional method of subjective refractometry with application of trial lenses was developed on the basis of data of large optical aberrations and significant depth of focus which values should be taken into account during interpretation of results of subjective refractometry. Our data regarding to prevalence of emmetropic refraction falls in line with basic science provisions in respect of the physiology of the eye.

© 2011 Spanish General Council of Optometry. Published by Elsevier España, S.L. All rights reserved.

PALABRAS CLAVE

Refractometría;
Profundidad de campo;
Profundidad de foco;
pseudacomodación

Método subjetivo de refractometría y profundidad de foco

Resumen

Objetivo: Estudiar el efecto de la profundidad de foco en la refracción subjetiva y la distribución de los estados refractivos miópicos e hipermetrópicos.

Métodos: Para el estudio se incluyeron 450 ojos de 305 sujetos con un intervalo de edad de 23 a 34 años. Se examinó la distribución de estados refractivos utilizando un método tradicional

* Corresponding author at: Eye Microsurgery Center, Komarov ave, 3, Kiev 03680, Ukraine.
E-mail address: nsmicro@gmail.com (N. Sergienko).




Figure 2 Displacement of the depth of focus in the course of subjective method of refractometry.

The purpose of this work was to study an impact of the depth of focus on subjective refraction and distribution of myopic and hyperopic refractive errors.

Methods

The subjects for this study were selected from the serving military officers of armed forces, Kiev. The medical records of about 3000 military men were reviewed to find persons with visual acuity of at least 1.0 or higher. The visual acuity was examined in 940 subjects, and then 450 eyes of 305 persons in the age range of 23–34 years were enrolled in the study.

All participants were aware of the purpose of the investigation and gave verbal consent.

The inclusion criteria were normal levels of visual acuity, a natural pupil diameter between 3 and 5 mm, no evidence of ocular diseases. The exclusion criteria were an astigmatism of $\geq 1.0D$, irregular astigmatism, any opacity of the ocular media which could impact a visual performance.

All the investigations were performed in the same room with uniform artificial illumination of about 210 lx. Visual acuity was tested at distance of 5 m using Landolt's broken ring chart fixed in the standard box having the local luminance about 145 cd/m². The lower 3 lines of the chart relate to a visual acuity 1.0, 1.5 and 2.0. These lines content the Landolt's circles with a size of 7.3 mm, 4.8 mm and 3.6 mm which at distance of 5 m subtend 5 min, 4.1 min and 2.5 min of arc, respectively.

A typical set of lenses was provided with spheres every quarter of a diopter to $\pm 4.0D$. Visual acuity scoring was performed if the subject correctly located Landolt's ring opening at least 5 times out of 7 attempts. The study was preceded by the preliminary screening examination of a

Table 1 Result of subjective refractometry using the traditional method.

Refraction	<i>n</i>	%
Hypermetropia	267	59.3
Myopia	99	22
Emmetropia	84	18.7

significant group of young persons in order to find eyes achieving visual acuity 2.0.

The degree of ametropia was assessed in two ways.

- (1) According to a traditional approach, the degree of hypermetropia was determined by the highest power convex lens and myopia by the lowest concave lens, which allowed the highest visual acuity to be reached: 1.0, 1.5 or 2.0. An eye was considered emmetropic if unaided visual acuity was at the highest level and adding a +0.25 D lens impaired acuity.
- (2) The data from hypermetropic eyes examined using the traditional method were corrected by a means of depth of focus. The degree of hypermetropia in an eye with visual acuity of 1.0, 1.5 or 2.0 was reduced at a depth of focus 1.0D, 0.7 D and 0.4D, respectively. A difference between the degree of hypermetropia and depth of focus less than 0.1 D was not taken into account.

Results

According to the traditional method of subjective refractometry, the distribution of refractions obtained in this study with the results of previous studies (Table 1). Hypermetropia occurred in 59% of cases and emmetropia in much fewer cases (18.4%).

As noted above, according to our concept, the depth of focus does not impact the incidence of myopia. Results regarding the degrees of hypermetropia and visual acuity are presented in Table 2. The eyes with light degrees of hypermetropia, +0.25 D or +0.5 D, were prevalent in 105 and 75 cases, respectively.

If we compare the hypermetropia +0.25 D with the depth of focus values (4 D, 0.7 D and 1.0 for visual acuity 2.0, 1.5 and 1.0, respectively), then these values are higher than the degree of hypermetropia. This finding was sufficient reason to consider all of these eyes as emmetropic. In a group of hypermetropia +0.5 D 4 eyes with visual acuity 2.0 had a depth of focus 0.4 D that was less than degree of hyperopia. These 4 eyes (Table 2) retained their place in the hypermetropic group. The other 71 eyes were changed to the emmetropic group.

The depth of focus of eyes with hypermetropia +0.75 D appeared to be equal to or greater than the degree of hypermetropia. Therefore, all 36 eyes were transferred to the emmetropic group. The depth of focus of eyes with hypermetropia +1.0 D and greater was also equal to or greater than the degree of hypermetropia. All of these eyes were regarded as hypermetropic.

As a result, the distribution of refractions changed (Table 3). The incidence of emmetropia rose to 65.3%, and

Table 2 Visual acuity and depth of focus of hyperopic eyes.

Degree of hypermetropia (diopters)	Visual acuity			Eyes changed to emmetropia	Eyes remained hypermetropic
	2.0	1.5	1.0		
+0.25	13	76	14	103	
+0.5	4	67	4	71	4
+0.75		36		36	
+1.0		23			23
Over +1.0		21	9		30
Depth of focus (diopters)	0.4	0.7	1.0	Total, 210	Total, 57

Table 3 Results of subjective refractometry on the basis of the depth of focus.

Refraction	<i>n</i>	%
Emmetropia	294	65.3
Myopia	99	22
Hypermetropia	57	12.7

hypermetropia occurred in only 12.7% of cases. The number of myopic eyes remained stable at 22%.

Discussion

Method of subjective refraction is associated with number of sources of uncertainty.¹² The depth-of-field/depth-of-focus are considered as the most important among them. The goal of this study was to clarify a role of the depth-of-focus in the subjective refractometry. We do not pretend to establish a new definition for refractive error or to claim a new distribution of refractive error in the general population. Our study cohort presents the pre-selected subjects with good and very good visual acuity. We ignored such factors as accommodation fluctuations and infinity vergence.

The real optical system of the human eye poses a number of the optical errors including aspheric shape refractive surfaces of the cornea and lens, decentration of the refractive surfaces, lens tilt, refractive irregularity inside the lens structure. All mentioned factors act simultaneously and summarize in the joint optical defect which now is called the wavefront aberration and can be measured by the aberrometers.

Aberrometry examination demonstrated that distribution of refractions within the optical zone varies between 1.0D and 1.5D. Emmetropic eyes are characterized by the both myopic and hyperopic refractions. The picture of distribution of refractions allows distinguishing the two components: irregular and regular astigmatism. The first is large and the second is too small.

The study of a relationship between the degree of wavefront aberrations and visual performance revealed a moderate correlation.¹³⁻¹⁶

The wavefront aberrations render a substantial impact on depth-of-focus and visual acuity. The depth-of-focus ± 0.35 D relates to visual acuity 1.5 which is close to an average value of visual acuity. The visual acuity 2.0 is characterized by

less ocular aberrations and shorter depth-of-focus ± 0.2 D. The eyes with the lowest limit of a normal vision 1.0 possess a significant amount of the aberrations and a depth of focus ± 0.55 D. The mentioned values were found for the pupil diameter 3 mm.

The aberration patterns are altering during an accommodation tension. This phenomenon does not relate to our study because all our measurements were carried out under condition of relaxed accommodation.

It is fundamental point how to consider in terms of refraction the eyes presented in Fig. 1A–C. When the retina is positioned in the middle of the depth-of-focus (Fig. 1A) the aberrometry map demonstrates irregular distribution of both myopic and hyperopic refractions with amplitude of 1.0–1.5 D.

In Fig. 1 anterior edge of the depth-of-focus touches the retina. Hyperopic refractions are dominant in the aberrometry map. At the same time the eye has normal visual acuity and optimal retinal image quality without help of accommodation. We believe that such eye should be considered as emmetropic.

When different degrees of myopia are prevalent in the aberrometry map (Fig. 1C), then however this eye is not myopic, but emmetropic because it possesses a normal unaided visual acuity.

The findings of our study have a more theoretical, rather than practical value.

Theoretically, emmetropia has always been considered the best for adapted refraction, and our data created a basis for verification of that point of view. Many papers regarding the prevalence of hypermetropia² have presented questions without proper answers. If nature selects for hypermetropia, then the reasons for this phenomenon should be determined. The necessity of constant accommodative tension, not only during near vision but also for far, appears to be a disadvantage of hypermetropia when compared with emmetropia. Our data on the prevalence of emmetropia make our knowledge about the distribution of refractions more consistent, and this distribution falls in line with basic science provisions regarding the physiology of the eye.

We do not insist on the implementation of our approach in the everyday practice of optometrists because the method requires more time, than conventional methods. Additionally, the depth of focus is a variable parameter that depends on pupil diameter, state of accommodation and pre-corneal tear film. It is impossible to know exactly the extent of the depth of focus, so approximations are inevitable.

However, there are practical implications that can be derived from this study. Namely, an estimation of preciseness of preoperative calculations of IOL optical power for the required target refraction requires postoperative refractometry. The refinement of the subjective method of refractometry does provide a guide for reconsidering the current methods. Light degrees of hypermetropia may not be regarded as an error. The majority of these eyes present emmetropia and predict the target refraction (Fig. 2).

References

1. Rosenfield M. Subjective refraction. In: *Optometry*. Elsevier; 2009. p. 209–28.
2. Duke-Elder S, Abrams D. *System of ophthalmology, vol. V. Variation of refractive state*. St. Louis: CV Mosby; 1970. p. 234–239.
3. Ciufreda KJ. Accommodation, the pupil, and presbyopia. In: Benjamin WJ, ed. *Borish's clinical refraction*. W.B. Saunders Company; 1998:77–120.
4. Oshima S. Studies on the depth-of-focus of the eye. *Jpn J Ophthalmol*. 1958;2:63–72.
5. von Bahr G. Studies of the depth-of-focus of the eye. *Acta Ophthalmol (Copenh)*. 1952;30:39–44.
6. Campbell FW. The depth of field of the human eye. *Opt Acta (Lond)*. 1957;4:157–164.
7. Charman WN, Whitefoot H. Pupil diameter and the depth-of-field of the human eye as measured by laser speckle. *Opt Acta*. 1977;24:1211–1216.
8. Tucker J, Charman WN. The depth-of-focus of the human eye for Snellen letters. *Am J Optom Physiol Opt*. 1975;53:2–3.
9. Ogle KN, Schwartz JT. Depth-of-focus of the human eye. *Am J Optom Physiol Opt*. 1959;49:273–280.
10. Sergienko NM, Tutchenko NN. Depth of focus: clinical manifestation. *Eur J Ophthalmol*. 2007;17:836–840.
11. Sergienko NM, Kondratenko YN, Tutchenko NN. Depth of focus in pseudophakic eyes. *Graefes Arch Clin Exp Ophthalmol*. 2008;246:1623–1627.
12. Smith G. Refraction and visual acuity measurements: what are measurement uncertainties? *Clin Exp Optom*. 2006;2:66–72.
13. Applegate RA, Ballentine Ch, Gross H, Sarver EJ, Sarver ChA. Visual acuity as function of Zernike mode and level of root mean square error. *Optom Vis Sci*. 2003;80:97–105.
14. Watson AB, Ahumada AJ. Predicting visual acuity from wave-front aberrations. *J Vis*. 2008;8:1–19.
15. Sergienko NM, Vladimirov DV, Molebny VV. Visual acuity and higher order aberrations. In: XXIV, *Congress of ESCRS*. 2006:155. Books of abstracts.
16. Sergienko NM, Vladimirov DV, Molebny VV. Influence of higher order aberrations on visual acuity. *Ophthalmol J*. 2009;5:14–17 [in Russian].